

MYCOTAXON

<http://dx.doi.org/10.5248/130.33>

Volume 130, pp. 33–40

January–March 2015

New records of one *Amygdalaria* and three *Porpidia* taxa (*Lecideaceae*) from China

LU-LU ZHANG, XIN ZHAO, & LING HU*

Key Laboratory of Plant Stress Research, College of Life Sciences, Shandong Normal University,
Jinan, 250014, P. R. China

* CORRESPONDENCE TO: lichenhuling@gmail.com

ABSTRACT —Four lichen taxa of *Lecideaceae*, *Amygdalaria consentiens* var. *consentiens*, *Porpidia carlottiana*, *P. lowiana*, and *P. tuberculosa*, are reported for the first time from China.

KEY WORDS —Asia, *Lecideales*, lichens

Introduction

The family *Lecideaceae* Chevall. contains about 23 genera and 547 species, of which the largest genus is *Lecidea* Ach., containing about 427 species (Kirk et al. 2008, Fryday & Hertel 2014). In China, twenty-three species of the other genera in *Lecideaceae* have been reported, including two each of *Amygdalaria* Norman, *Bellemerea* Hafellner & Cl. Roux, and *Immersaria* Rambold & Pietschm.; one each of *Lecidoma* Gotth. Schneid. & Hertel, *Paraporpidia* Rambold & Pietschm., and *Stenhammarella* Hertel; and 16 species of *Porpidia* Körb. (Hertel 1977, Wei 1991; Aptroot & Seaward 1999; Aptroot 2002; Aptroot & Sparrius 2003; Obermayer 2004; Guo 2005; Zhang et al. 2010, 2012; Wang et al. 2012; Ismayi & Abbas 2013; Hu et al. 2014).

Amygdalaria and *Porpidia* are obviously very closely related. Both have large halonate ascospores, a high hymenium, *Porpidia*-type asci, and a dark pigmented hypothecium and are (with a few exceptions) restricted to lime-free, silicate rocks. However *Amygdalaria* can be best distinguished from *Porpidia* by the presence of cephalodia, the higher hymenium (over 130 μm), the larger ascospores (generally $20\text{--}35 \times 10\text{--}16 \mu\text{m}$) with conspicuous, rather compact episporos, and a tendency toward a brownish or yellowish pink thallus (Inoue 1984, Brodo & Hertel 1987, Gowan 1989, Smith et al. 2009).

During our research on the lecideoid taxa of *Lecideaceae* in China, we have identified four taxa new to the country: *Amygdalaria consentiens* var. *consentiens*, *Porpidia carlottiana*, *P. lowiana*, and *P. tuberculosa*.

Materials & methods

The studied specimens are preserved in the Lichen Section of Botanical Herbarium, Shandong Normal University, Jinan, China (SDNU); Herbarium Mycologicum Academiae Sinicae – Lichenes, Beijing, China (HMAS-L); and the Herbarium of the Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China (KUN). The morphological and anatomical characters were examined under a stereomicroscope (COIC XTL7045B2) and a polarizing microscope (OLYMPUS CX41). The thalli and medullae were tested for identification with K (10% aqueous KOH solution), C (saturated aqueous NaClO solution), I (a 10% aqueous KI solution), P (a saturated *p*-phenylenediamine solution in 95% ethyl alcohol). The lichen substances were identified using standardized thin layer chromatography techniques (TLC) with solvent system C (Orange et al. 2010). Photos of these lichens were taken under OLYMPUS SZX16 and BX61 with DP72.

Taxonomic descriptions

Amygdalaria consentiens (Nyl.) Hertel, Brodo & Mas. Inoue, J. Hattori Bot. Lab. 56: 322 (1984) var. *consentiens* FIG. 1

MORPHOLOGY — **THALLUS** continuous, cracked-areolate, usually thick, light orange to brownish gray, often becoming oxidated orange; medulla I–; prothallus black, thin and conspicuous; cephalodia mostly flush with thallus surface, almost the same colour as the thallus, but occasionally pinker or gray, containing *Stigonema*; soredia absent. **APOTHECIA** entirely sunken in thallus, (0.35–)0.45–0.9 mm diam., usually one or two per areole; disc dark brown to black, margin slightly to moderately rusty orange pruinose. **EXCIPLE** black-brown, 50–70 µm wide, without crystals; epihymenium brownish, with orange crystals; hymenium hyaline, 150–200 µm tall; subhymenium colorless, 45–60 µm tall; hypothecium blackish brown, usually thin, bowl-shaped; paraphyses strongly anastomosed and branched, almost moniliform at the apical part, with ±swollen apices. **ASCI** clavate, *Porpidia*-type; ascospores hyaline, simple, ellipsoid, 28–32(–35) × 14–17(–19) µm, halonate. **PYCNIDIA** immersed, conidia bacilliform, 6–9 × c. 1 µm.

CHEMISTRY — Thallus and medulla K–, C–, KC–, P–. No lichen substances were detected by TLC.

DISTRIBUTION — *Amygdalaria consentiens* has been reported from Asia, Europe, and North America (Inoue 1984, Brodo & Hertel 1987, Smith et al. 2009). New to China.

SPECIMEN EXAMINED: CHINA. YUNNAN, Luquan, Mt. Jiaozixueshan, alt. 3800 m, on rock, 26 Oct. 2008, Z.J. Ren 20108488 (SDNU).




FIG. 1 *Amygdalaria consentiens* var. *consentiens* (Ren 20108488, SDNU). A: thallus; B: prothallus; C: apothecium section; D: epithymenium with crystals; E: amyloid reaction of ascus; F: ascus and ascospores; G: ascospores; H: paraphyses.

COMMENTS — *Amygdalaria consentiens* var. *consentiens* is morphologically similar to *A. consentiens* var. *japonica* and *A. continua*, but *A. consentiens* var. *japonica* contains stictic and constictic acids, while *A. continua* has a smooth to finely rimose thallus which is rarely cracked-areolate.

Porpidia carlottiana Gowan, Bryologist 92: 39 (1989)

FIG. 2

MORPHOLOGY — THALLUS crustose, finely cracked to subrimose, 0.15–0.25 mm thick, whitish gray; medulla I–; prothallus continuous between thallus




FIG. 2 *Porpidia carlottiana* (Wang 20127123, SDNU). A: thallus; B: apothecium section; C: amyloid reaction of ascus; D: ascospores; E: exciple and epihymenium without crystals.

patches, black, thin; soredia absent. APOTHECIA scattered or contiguous, sunken in thallus when young, becoming sessile on a broad base when mature, 0.5–1.1(–1.5) mm diam.; disc black, plane to weakly convex, usually with heavy whitish pruina; margin bare, distinct, regular to flexuose. EXCIPLE dark brown to black at margin, brown within, 90–105 µm wide, without crystals; epihymenium yellowish brown or olive brown, without crystals; hymenium hyaline, 90–105(–130) µm tall; subhymenium 20–25 µm, hypothecium blackish brown; paraphyses strongly anastomosed and apically branched. ASCI clavate, *Porpidia*-type; ascospores hyaline, simple, ellipsoid, 17–20 × 7–10 µm, halonate. PYCNIDIA not observed.

CHEMISTRY — Thallus and medulla K–, C–, KC–, P–. 2'-O-methyl-superphyllinic acid was detected by TLC.

DISTRIBUTION — *Porpidia carlottiana* has been reported from North America (Gowan 1989). New to China.

SPECIMEN EXAMINED: CHINA. YUNNAN, Lijiang, Mt. Laojunshan, alt. 3800 m, on rock, 5 Nov. 2009, H.Y. Wang 20127123 (SDNU).

COMMENTS — *Porpidia carlottiana* is closely related to *P. rugosa*, which also produces pruinose apothecia and 2'-O-methylsuperphyllinic acid. However, *P. rugosa* always has soredia, and the apothecia rarely seen.

Porpidia lowiana Gowan, Bryologist 92: 49 (1989)

FIG. 3

MORPHOLOGY — **THALLUS** crustose, continuous, finely cracked to rimose-areolate, 0.2–0.5 mm thick, yellowish gray to whitish, sometimes dark gray or greenish gray; medulla I–; prothallus absent; soredia absent. **APOTHECIA** scattered to contiguous, soon becoming broadly sessile, 0.5–1.3 mm diam.; disc black, non-pruinose to moderately pruinose; margin bare, distinct, even to weakly crenulate, regular to flexuose. **EXCIPLE** mostly black and carbonaceous at exciple margin, hyaline within, 55–90 μm wide, with crystals that are soluble in K and C; epihymenium olive brown, without crystals; hymenium hyaline, 70–110 μm tall; subhymenium 18–26 μm , hypothecium blackish brown; paraphyses strongly anastomosed and apically branched. **ASCI** clavate, *Porpidia*-type; ascospores hyaline, simple, ellipsoid or tear-drop-shape, 14–16(–25) $\times$ (4.5–)6–8 μm , halonate. **PYCNIIDIA** not observed.

CHEMISTRY — Thallus and medulla K–, C–, KC–, P–. Confluent, 2'-O-methylmicrophyllinic, and 2'-O-methylperlatolic acids were detected by TLC.

DISTRIBUTION — *Porpidia lowiana* has been reported from Europe and North America (Gowan 1989, Smith et al. 2009). New to China.




FIG. 3 *Porpidia lowiana* (Ren 20102097, SDNU). A: thallus; B: apothecium section; C: amyloid reaction of ascus; D: ascospores; E: exciple with crystals.

SPECIMENS EXAMINED: CHINA. HELONGJIANG, Tahe, Mt. Mengkeshan, alt. 555 m, on rock, 14 Aug. 2009, Q. Ren 20102097 (SDNU). INNER MONGOLIA, Mangui, alt. 900 m, on rock, 13 Sep. 1977, J.C. Wei, 074581 (HMAS-L); SICHUAN, Dukou, Mt. Dabaoding, alt. 1950 m, on rock, 21 Jun. 1983, L.S. Wang 83-184 (KUN); YUNNAN, Xiping, Mt. Mopan, alt. 2000 m, on rock, 20 Dec. 2008, L.S. Wang 08-30003 (KUN); Chuxiong, Mt. Zixi, alt. 2060 m, on rock, 31 Aug. 2005, L.S. Wang 05-25263 (KUN); Lincang county, Matai village, alt. 1650 m, on rock, 20 Oct. 2003, L.S. Wang 03-22921, 03-22922 (KUN); Yimen, alt. 1600 m, on rock, 10 Dec. 1990, L.S. Wang 11707 (KUN).

COMMENTS — *Porpidia lowiana* is closely related to *P. cinereoatra*, from which it differs mainly in having a thinner, yellowish thallus, thicker proper margin (c. 0.05 mm in *P. cinereoatra*), and sessile apothecia (mainly sunken in *P. cinereoatra*).

Porpidia tuberculosa (Sm.) Hertel & Knoph, Beih. Nova Hedwigia 79: 438 (1984)

FIG. 4

MORPHOLOGY — THALLUS crustose, continuous, subrimose to distinctly rimose-areolate, 0.2–0.3(–0.5) mm thick, medium gray or sometimes rusty orange; medulla I+ violet; prothallus usually present when abutting another lichen, black; soralia present, 0.3–0.7 mm diam., round to irregular, scattered to crowded, shallowly erose or abraded, usually with slightly raised rim; soredia farinose to granular, white, pale gray or flecked blue-gray or blackish, I+ violet. APOTHECIA occasional, scattered, soon becoming sessile, 0.9–1.5 (–1.7) mm diam.; disc black, plane to subconvex, weakly to moderately pruinose; margin bare, distinct, regular to flexuose. EXCIPLE greenish to brownish black at exciple margin, dark brown within, 90–140 µm wide, without crystals; epihymenium olive brown, without crystals; hymenium hyaline, 90–105 µm tall; subhymenium 35–50 µm, hypothecium blackish brown; paraphyses strongly anastomosed and apically branched. ASCI clavate, *Porpidia*-type; ascospores hyaline, simple, ellipsoid, $17.5\text{--}22 \times 7\text{--}8$ µm, halonate. PYCNIDIA immersed, conidia bacilliform, $7.5\text{--}10 \times c. 1$ µm.

CHEMISTRY — Thallus and medulla K–, C–, KC–, P–. Confluent, 2'-O-methylmicrophyllinic, and 2'-O-methylperlatolic acids were detected by TLC.

DISTRIBUTION — *Porpidia tuberculosa* has been reported from Asia, Macaronesia, Europe, and North & South America (Hertel 1984, Smith et al. 2009). New to China.

SPECIMENS EXAMINED: CHINA. HUBEI, Shennongjia, Mt. Dashennongjia, alt. 3000 m, on rock, 27 Jun. 1984, J.C. Wei, 067994, 068005, 074292 (HMAS-L). YUNNAN, Luquan, Mt. Jiaozixueshan, alt. 3800 m, on rock, 27 Oct. 2008, H.Y. Wang 20081180, 20082302 (SDNU).

COMMENTS — *Porpidia tuberculosa*, *P. rugosa* and *P. soredizodes* all have a thallus with soralia. However, *P. rugosa* has a more verrucose thallus,




FIG. 4 *Porpidia tuberculosa* (Wei 067994, HMAS-L). A: thallus; B: apothecium section; C: amyloid reaction of ascus; D: conidia; E: exciple and epihymenium without crystals.

I- medulla, and contains 2'-O-methylsuperphyllinic and glaucophaeic acids. *Porpidia soredizodes* has an I- medulla as well as an overall smaller thinner (≤ 0.2 mm) darker thallus that contains stictic acid.

Acknowledgements

The authors thank Dr. A. Aptroot (ABL Herbarium, Soest, the Netherlands) and Dr. Shou-Yu Guo (State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China) for presubmission reviews. This study was supported by Program for Scientific Research Innovation Team in Colleges and Universities of Shandong Province, the National Natural Science Foundation of China (31170187), Foundation of Key Laboratory, CAS (KLBB-201306), and the Scientific Research Foundation of Graduate School of Shandong Normal University (BCX1406).

Literature cited

- Aptroot A. 2002. Corticolous and saxicolous lichens from Xishuangbanna, southern Yunnan, China. <http://www.nhm.uio.no/botanisk/lav/Yunnan>.
- Aptroot A, Seaward MRD. 1999. Annotated checklist of Hong Kong lichens. *Tropical Bryology* 17: 57–101.
- Aptroot A, Sparrius LB. 2003. New microlichens from Taiwan. *Fungal Diversity* 14: 1–50.

- Brodo IM, Hertel H. 1987. The lichen genus *Amygdalaria* (*Porpidiaceae*) in North America. *Herzogia* 7: 493–521
- Fryday AM, Hertel H. 2014. A contribution to the family *Lecideaceae* s. lat. (*Lecanoromycetidae* inc. sed., lichenized *Ascomycota*) in the southern subpolar region; including eight new species and some revised generic circumscriptions. *Lichenologist* 46(3): 389–412.
<http://dx.doi.org/10.1017/S0024282913000704>
- Gowan SP. 1989. The lichen genus *Porpidia* (*Porpidiaceae*) in North America. *Bryologist* 92: 25–59.
- Guo SY. 2005. Lichens. 31–82, in: WY Zhuang (ed.). *Fungi of northwestern China*. Mycotaxon Ltd., Ithaca, New York.
- Hertel H. 1977. Gesteinsbewohnende Arten der Sammelgattung *Lecidea* (*Lichenes*) aus Zentral-, Ost- und Südasien. Khumbu Himal, Ergebnisse des Forschungsunternehmens Nepal Himalaya. 6: 145–378.
- Hertel H. 1984. Über saxicole, lecideoide Flechten der Subantarktis. Beihefte zur Nova Hedwigia. 79: 399–499.
- Hu L, Zhao X, Sun LY, Zhao ZT, Zhang LL. 2014. Four lecideoid lichens new to China. *Mycotaxon* 128: 83–91. <http://dx.doi.org/10.5248/128>
- Inoue M. 1984. Japanese crustose lichen genera formerly reported under *Lecidea sensu lato*. 1. *Amygdalaria* Norman. *Journal of the Hattori Botanical Laboratory* 56: 321–330.
- Ismayi G, Abbas A. 2013. *Bellemerea alpina* — new lichen species record from China. *Plant Science Journal* 31(4): 333–335.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. *Dictionary of the fungi*. 10th edition. CABI Bioscience: CAB International. 771 p.
- Obermayer W. 2004. Additions to the lichen flora of the Tibetan region. *Bibliotheca Lichenologica* 88: 479–526.
- Orange A, James PW, White FJ. 2010. *Microchemical methods for the identification of lichens*. 2nd edition. London: British Lichen Society.
- Smith CW, Aptroot A, Coppins BJ, Fletcher A, Gilbert OL, James PW, Wolseley PA (eds). 2009. *The lichens of Great Britain and Ireland*. Natural History Museum Publications, in association with The British Lichen Society.
- Wang XY, Zhang LL, Joshi Y, Wang HY, Hur JS. 2012. New species and new records of the lichen genus *Porpidia* (*Lecideaceae*) from western China. *Lichenologist* 44(5): 619–624.
<http://dx.doi.org/10.1017/S0024282912000242>
- Wei JC. 1991. *An enumeration of lichens in China*. International Academic Publishers, Beijing.
- Zhang LL, Wang HY, Sun LY, Zhao ZT. 2010. Four lichens of the genus *Lecidea* from China. *Mycotaxon* 112: 445–450. <http://dx.doi.org/10.5248/112.445>
- Zhang LL, Wang LS, Wang HY, Zhao ZT. 2012. Four new records of lecideoid lichens from China. *Mycotaxon* 119: 445–451. <http://dx.doi.org/10.5248/119.445>